tebusha

Oil in water emulsion containing 250 g/L tebuconazole.
Broad spectrum fungicide for use in winter and spring crops of wheat (excluding durum), barley, oats, rye, triticale and oilseed rape.

SAFETY INFORMATION

DANGER
May cause an allergic skin reaction
Causes serious eye damage
Suspected of damaging the unborn child
Toxic to aquatic life with long lasting effects

Keep out of reach of children.
Do not eat, drink or smoke when using this product.
Avoid breathing vapours/spray.
Wear protective gloves/protective clothing/eye protection.
IF ON SKIN: Wash with plenty of soap and water.
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
If skin irritation or rash occurs. Get medical advice/attention.
Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste
To protect aquatic organisms respect an unsprayed buffer zone of 5 m to surface water bodies.
To avoid risks to human health and the environment comply with the instructions for use.

PCS No. 04884

Batch number: see container
Registration holder: Sharda Europe b.v.b.a., Jozef Mertensstraat 142, 1702 Dilbeek, Belgium
Distributed by: Sharda Europe b.v.b.a., Jozef Mertensstraat 142, 1702 Dilbeek, Belgium. Phone no: +91 22 5678 2800

Emergency number: +91 22 5678 2800 (8.00-16.00h)

Sharda Cropchem

SHARDA CROPCHEM LIMITED
(previously Sharda Worldwide Exports Pvt. Ltd)
Prime Business Park, Dashrathlal Joshi Road
Vile Parle (West) Mumbai - 400056 , India
Tel : + 91 22 6678 2800; Fax : + 91 22 6678 2828

UFI: KARA-26QR-U70V-2WDN

PRECAUTIONS
If you feel unwell seek medical advice immediately (show the label where possible).

In case of accident, call the National Poisons Information Centre:
Tel: +353 (0)1 809 2566 or +353 (0)1 837 9964
Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads. Protect from frost.

READ LABEL CAREFULLY BEFORE USE

IMPORTANT INFORMATION			
FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE			
Crops:	**Maximum individual dose:**	**Maximum total dose:**	**Latest time of application:**
Cereals (winter & spring wheat, winter & spring barley, oats, triticale and rye)	1 L/ha	2 L/ha	Before grain watery ripe stage (GS 71)
Oilseed rape (winter and spring)	1 L/ha	2 L/ha	Before full flowering (GS 65)

Tebusha is a fungicide for the control of a wide range of fungal diseases.

INSTRUCTIONS FOR USE

For best disease control and yield, apply the product at an early stage of disease development, before infection spreads to new growth

To avoid the occurrence of resistance, do not apply the product or similar products with the azole-type active substance other than for preventive reasons or as soon as possible after the occurrence of infection. Do not rely on curative potential of this type of active ingredient.

To avoid occurrence of resistance, do not apply the product or similar products based exclusively on azole-type active ingredient in the crop consecutively without interruption by treatment using a different fungicide with a different mechanism of efficiency.

Avoid drift onto plants outside the target area due to risk of damage.

Diseases Controlled-

Wheat: Reduction of *Septoria* leaf blotch, Glume blotch and *Fusarium* ear blight. Control of yellow rust, brown rust and powdery mildew.
Barley: Reduction of *Fusarium* ear blight. Control of brown rust and powdery mildew.
Oat: Control of powdery mildew and crown rust.
Triticale: Reduction of *Septoria* leaf blotch and Glume blotch. Control of yellow rust, brown rust and powdery mildew.
Rye: Control of yellow rust, brown rust and powdery mildew.
Oilseed rape: Control of *Alternaria*, *Sclerotinia* and *Phoma* leaf spot.

Tebusha acts as a growth regulator in oilseed rape.

Crop sensitivity: For advice on crop sensitivity and replacement after crop failure, consult with distributor.

Resistance
To prevent resistance always follow label recommendations and monitor fields regularly for any signs of poor control. Use fungicides with different modes of action.

CROP SPECIFIC RECOMMENDATIONS

Wheat, barley, rye, triticale, oats
Apply at first sign of disease up until before grain watery ripe stage of growth. A second application may be required two to three weeks later.
Water volumes
Most crops: 100 – 200 L/ha water
Dense crops after first node detectable stage: 250 L/ha water
Larger crops and varieties highly susceptible to disease: 300 L/ha water

Oilseed rape: Apply in 100 – 400 l/ha water at any time up to and including the full flowering stage.

MIXING, APPLICATION AND CLEANING
Never prepare more spray solution than required.
Thoroughly shake pack before use.
When preparing the mixture, half fill the tank with clean water and start the agitation.
Add Tebusha in the tank under agitation and then fill the tank with water. Continue agitation until spraying is completed. Apply using a tractor mounted/ trailed sprayer.

Use a medium quality spray to ensure a good and even spray cover of the target. A spray pressure of 2-3 bar is recommended.

Avoid spillage. In case of spill, remove contaminated clothes and wash skin immediately.

Clean equipment immediately after use with a spray tank cleaner.

Empty container and rinse with an integrated pressure rinsing device or manually rinsing three times.